

JACOBI'S TRIPLE PRODUCT, MOCK THETA FUNCTIONS, UNIMODAL SEQUENCES AND THE q -BRACKET

ROBERT SCHNEIDER

ABSTRACT. In Ramanujan's final letter to Hardy, he wrote of a strange new class of infinite series he called "mock theta functions". It turns out all of Ramanujan's mock theta functions are essentially specializations of a so-called universal mock theta function $g_3(z, q)$ of Gordon–McIntosh. Here we show that g_3 arises naturally from the reciprocal of the classical Jacobi triple product—and is intimately tied to rank generating functions for unimodal sequences, which are connected to mock modular and quantum modular forms—under the action of an operator related to statistical physics and partition theory, the q -bracket of Bloch–Okounkov. Secondly, we find $g_3(z, q)$ to extend in the variable q to the entire complex plane minus the unit circle, and give a finite formula for this universal mock theta function at roots of unity, that is simple by comparison to other such formulas in the literature; we also indicate similar formulas for other q -hypergeometric series. Finally, we look at interesting "quantum" behaviors of mock theta functions inside, outside, and on the unit circle.

1. INTRODUCTION: SEA CHANGE

When Ramanujan put to sea from India for Cambridge University in 1914, destined to revolutionize number theory, a revolution in physics was already full-sail in Europe. Just one year earlier, the Rutherford–Bohr model of atomic shells heralded the emergence of a paradoxical new *quantum theory* of nature that contradicted common sense. In 1915, Einstein would describe how space, light, matter, geometry itself, warp and bend in harmonious interplay. The following year, Schwarzschild found Einstein's equations to predict the existence of monstrously inharmonious *black holes*, that in fact we can study directly (just very recently) using interstellar gravitational waves [1] as well as in Earth-bound laboratories [41].

During Ramanujan's five years working with G. H. Hardy, news of the paradigm shift in physics must have created a thrill among the mathematicians at Trinity College, Isaac Newton's *alma mater*. Certainly Hardy would have been aware of the sea change. After all, J. J. Thomson's discovery of the electron, as well as his subsequent "plum-pudding" atomic model, had been made at Cambridge's Cavendish Laboratory; Rutherford had done his post-doctoral work with Thomson there; and Bohr came to Cambridge to work under Thomson in 1911 [20]. Moreover, Hardy's intellectual colleague Hilbert was in a public race with Einstein to write down the equations of General Relativity [25].

We don't know how aware Ramanujan was of these happenings in physics, yet his flights of imagination and break with academic tradition were expressions of the scientific *Zeitgeist* of the age. After returning to India in 1919, as he approached his own tragic event horizon, Ramanujan's thoughts ventured into realms that—like the domains of

This material is based upon work partially supported by the National Science Foundation under grant DMS-1128155.

subatomic particles and gravitational waves—would require the technology of a future era to navigate [33].

In the final letter he sent to Hardy, dated 1920, Ramanujan described a new class of mathematical objects he called *mock theta functions* [34], that mimic certain behaviors of classical *modular forms* [6, 32]. These q -series turn out to have profoundly curious analytic, combinatorial and algebraic properties. Ramanujan gave a prototypical example $f(q)$ of a mock theta function, defined by the q -series

$$(1) \quad f(q) := \sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q; q)_n^2}.$$

He claimed that $f(q)$ is “almost” modular in a number of ways. For instance, Ramanujan provided a pair of functions $\pm b(q)$ with

$$b(q) := (q; q^2)_{\infty} (1 - 2q + 2q^4 - \dots)$$

that are modular up to multiplication by $q^{-1/24}$, to compensate for the singularities arising in the denominator of (1) as q approaches an even-order root of unity ζ_{2k} radially from within the unit circle:

$$(2) \quad \lim_{q \rightarrow \zeta_{2k}} (f(q) - (-1)^k b(q)) = \mathcal{O}(1)$$

This was confirmed by Watson in 1936 [43], and quantifies to some degree just how “almost” modular $f(q)$ is: at least at even-order roots of unity, $f(q)$ looks like a modular form plus a constant.

Only in the twenty-first century have we begun to grasp the larger meaning of functions such as this, in the wake of Zwegers’s innovative Ph.D. thesis of 2002 [31, 47]. We now know Ramanujan’s mock theta functions are examples of *mock modular forms*, which are the holomorphic parts of more complicated creatures called *harmonic Maass forms* [31]. In 2012, Folsom–Ono–Rhoades [19] made explicit the limit given in (2), showing that

$$(3) \quad \lim_{q \rightarrow \zeta_{2k}} (f(q) - (-1)^k b(q)) = -4U(-1, \zeta_{2k}),$$

where $U(z, q)$ is the *rank generating function for strongly unimodal sequences* in combinatorics (see [9]), and is closely related to *partial theta functions* and mock modular forms. By this connection to U , the work of Folsom–Ono–Rhoades along with Bryson–Ono–Pitman–Rhoades [11] reveals that the mock theta function $f(q)$ is strongly connected to the newly-discovered species of *quantum modular forms* in the sense of Zagier: functions that are modular up to the addition of some “suitably nice” function, and (in Zagier’s words) have “the ‘feel’ of the objects occurring in perturbative quantum field theory” [46].

We do not know what sparked Ramanujan to discover mock theta functions, but we see in this paper that they are indeed natural forms to study from a classical perspective. It turns out in Section 2 that all of his mock theta functions—to be precise, the odd-order *universal mock theta function* of Gordon–McIntosh that specializes to the mock theta functions Ramanujan wrote down [21]—arise from the *Jacobi triple product*, a fundamental object in number theory and combinatorics [7], and are generally “entangled” with rank generating functions for unimodal sequences, under the action of the q -*bracket* operator from statistical physics and partition theory that boils down to multiplication by $(q; q)_{\infty}$ [45]. In Section 3 we find finite formulas for the odd-order universal mock theta function

and indicate similar formulas for other q -hypergeometric series. In some cases, we do encounter the “feel” of quantum theory.

2. CONNECTING THE TRIPLE PRODUCT TO MOCK THETA FUNCTIONS

At the wildest boundaries of nature, we see tantalizing hints of q -series. Recall that Borchers proposed a proof of the Jacobi triple product formula based on the *Dirac sea* model of the quantum vacuum (see [13]); it is as if this beautiful, versatile identity emerges from properties of empty space. Also from the universe of q -hypergeometric series, mock theta functions and their generalization mock modular forms [31] are connected conjecturally to deep mysteries in physics, like unthinkable phenomena at the edges of black holes [16, 15]. All the diversity of physical reality—and of our own mental experience—plays out quite organically between these enigmatic extremes. Perhaps not unrelatedly, in this study we see there is an organic connection between the Jacobi triple product and mock theta functions.

For brevity, we refer the reader to the author’s related paper [39] for standard notations and terminology, and move directly to introducing the principal actors. We take $q, z \in \mathbb{C}$, $|q| < 1$, $z \neq 0$ throughout, unless stated otherwise.

The *Jacobi triple product* $j(z; q)$ is implicit in countless famous classical identities (see [7]). Up to multiplication by rational powers of q , the triple product specializes to the Jacobi theta function (that Ramanujan constructed “mock” versions of), a weight $1/2$ modular form which is also important in physics as a solution to the heat equation. The triple product is defined by

$$(4) \quad j(z; q) := (z; q)_\infty (z^{-1}q; q)_\infty (q; q)_\infty,$$

where $|q| < 1$ and $(z; q)_0 := 1$, $(z; q)_n := \prod_{0 \leq i \leq n-1} (1 - zq^i)$, $(z; q)_\infty := \lim_{n \rightarrow \infty} (z; q)_n$ is the usual q -Pochhammer symbol. We note the simple zero at $z = 1$ from the $(1 - z)$ factor in $(z; q)_\infty$.

The odd-order *universal mock theta function* $g_3(z, q)$ of Gordon and McIntosh [21], which specializes to Ramanujan’s mock theta functions up to changes of variables and multiplication by rational powers of q , is defined as

$$(5) \quad g_3(z, q) := \sum_{n=1}^{\infty} \frac{q^{n(n-1)}}{(z; q)_n (z^{-1}q; q)_n},$$

and, like the triple product, is subject to all sorts of wonderful transformations. We note the simple pole at $z = 1$.

Let us recall that a *unimodal sequence* of integers is of the type

$$0 \leq a_1 \leq a_2 \leq \dots \leq a_r \leq \bar{c} \geq b_1 \geq b_2 \geq \dots \geq b_s \geq 0.$$

The term $\bar{c}$ is called the *peak* of the sequence; generalizing this concept, if $\bar{c}$ occurs with multiplicity $\geq k$, we might consider the unimodal sequence with a k -fold *peak*

$$0 \leq a_1 \leq a_2 \leq \dots \leq a_r \leq \bar{c} \bar{c} \dots \bar{c} \geq b_1 \geq b_2 \geq \dots \geq b_s \geq 0,$$

where “ $\bar{c} \bar{c} \dots \bar{c}$ ” denotes k repetitions of $\bar{c}$. When all the inequalities above are strictly “ $<$ ” or “ $>$ ” the sequence is *strongly unimodal*.

If r is the number of a_i to the left and s is the number of b_j to the right of a unimodal sequence, the difference $s - r$ is called the *rank* of the sequence. Another series that plays a role here is the *rank generating function* $\tilde{U}(z, q)$ for unimodal sequences, given by

$$(6) \quad \tilde{U}(z, q) := \sum_{n=0}^{\infty} \frac{q^n}{(zq; q)_n (z^{-1}q; q)_n},$$

where $(z^{-1}q; q)_n$ in the denominator gives rise to the a_i , and $(zq; q)_n$ generates the b_j (see [11]). If we replace z with $-z$, the right-most series is actually the very first expression Andrews revealed from Ramanujan's "lost" notebook ([4], Eq. 1.1) shortly after unearthing the papers at Trinity College [36]. This form, which is related to partial theta functions [27], was swimming alongside mock theta functions in the Indian mathematician's imagination during his final year.

Finally, following Bloch–Okounkov [8] as well as Zagier [45], we define the q -*bracket* $\langle f \rangle_q$ of a function $f : \mathcal{P} \rightarrow \mathbb{C}$, with $\mathcal{P}$ the set of integer partitions, to be given by

$$(7) \quad \langle f \rangle_q := \frac{\sum_{\lambda \in \mathcal{P}} f(\lambda) q^{|\lambda|}}{\sum_{\lambda \in \mathcal{P}} q^{|\lambda|}} = (q; q)_{\infty} \sum_{\lambda \in \mathcal{P}} f(\lambda) q^{|\lambda|},$$

where the sums are taken over partitions λ , with $|\lambda|$ the *size* of a partition (i.e., the number being partitioned). This operator represents the expected value in statistical physics of a measurement over a grand ensemble whose states are indexed by partitions with weights f , for a canonical choice of q ; this is the meaning of the quotient in the middle of (7).

However, we proceed formally here using the right-most expression, without drawing too much physical interpretation (while always keeping the mysterious feeling that our formulas resonate in physical reality). Simply multiplying by $(q; q)_{\infty}$ induces quite interesting q -series phenomena: Bloch–Okounkov [8], Zagier [45], and Griffin–Jameson–Trebart-Leder [22] show that the q -bracket can produce families of modular, quasimodular and p -adic modular forms; and the present author finds the q -bracket to play a natural role in partition theory as well [39], modularity aside. (We highly recommend Zagier's paper [45] for more about the q -bracket.)

It turns out that the reciprocal of the Jacobi triple product

$$j(z; q)^{-1} =: \sum_{\lambda \in \mathcal{P}} j_z(\lambda) q^{|\lambda|}$$

has a very rich and interesting interpretation in terms of the q -bracket operator, which in this case has the shape

$$\langle j_z \rangle_q = (q; q)_{\infty} j(z; q)^{-1} = \frac{1}{(z; q)_{\infty} (z^{-1}q; q)_{\infty}}.$$

Note that this q -bracket also has a simple pole at $z = 1$. We abuse notations somewhat in writing the coefficients j_z in this way, as if $z \in \mathbb{C}$ were a constant. In fact, j_z is a map from the partitions to $\mathbb{Z}[z]$, which the author gives explicitly in [39] as

$$(8) \quad j_z(\lambda) = (1 - z)^{-1} \sum_{\delta | \lambda} \sum_{\varepsilon | \delta} z^{\text{crk}(\varepsilon)}$$

for $z \neq 1$, where " $\alpha | \beta$ " means $\alpha \in \mathcal{P}$ is a subpartition of $\beta \in \mathcal{P}$, and "crk" denotes the *crank* statistic of Andrews–Garvan [5] defined below.

Definition 1. The crank $\text{crk}(\lambda)$ of a partition λ is equal to its largest part if the multiplicity $m_1(\lambda)$ of 1 as a part of λ is 0 (that is, there are no 1's), and if $m_1(\lambda) > 0$ then $\text{crk}(\lambda) = \#\{\text{parts of } \lambda \text{ that are larger than } m_1(\lambda)\} - m_1(\lambda)$.

Remark. The well-known crank generating function (see [19]) is given by

$$C(z; q) = \frac{(q; q)_\infty}{(zq; q)_\infty (z^{-1}q; q)_\infty} = (1 - z)(q; q)_\infty \langle j_z \rangle_q.$$

Expanding $C(z; q)$ as a power series in q , the n th coefficient is given by $\sum_{\lambda \vdash n} z^{\text{crk}(\lambda)}$, where “ $\lambda \vdash n$ ” means λ is a partition of n .

In [39] we used the q -bracket operator to find the coefficients of $\langle j_z \rangle_q$ explicitly in terms of sums over subpartitions and the crank statistic, as well. Now we take a different approach, and look at $\langle j_z \rangle_q$ from the point-of-view of q -hypergeometric relations. It turns out the odd-order universal mock theta function g_3 (in an “inverted” form) and the unimodal rank generating function $\tilde{U}$ naturally arise together as components of $\langle j_z \rangle_q$.

Theorem 2. For $0 < |q| < 1, z \neq 0, z \neq 1$, the following statements are true:

(i) We have the q -bracket formula

$$\langle j_z \rangle_q = 1 + [z(1 - q) + z^{-1}q] g_3(z^{-1}, q^{-1}) + \frac{zq^2}{1 - z} \tilde{U}(z, q).$$

(ii) The “inverted” mock theta function component in part (i) converges, and can be written in the form

$$g_3(z^{-1}, q^{-1}) = \sum_{n=1}^{\infty} \frac{q^n}{(z; q)_n (z^{-1}q; q)_n}.$$

By considering the factor $z(1 - q) + z^{-1}q$ as $|z| \rightarrow \infty$ and as $|z| \rightarrow 0$ in part (i) of Theorem 2, we get the following asymptotics.

Corollary 3. We have the asymptotic estimates:

(i) For $0 < |q| < 1 \ll |z|$, we have

$$\langle j_z \rangle_q \sim z(1 - q)g_3(z^{-1}, q^{-1}) \text{ as } |z| \rightarrow \infty.$$

(ii) For $0 < |q| < 1, 0 < |z| \ll 1$, we have

$$\langle j_z \rangle_q \sim z^{-1}q g_3(z^{-1}, q^{-1}) \text{ as } |z| \rightarrow 0.$$

Thus the inverted mock theta function component dominates the behavior of the q -bracket for z not close to the unit circle (which is “most” of the complex plane).

Remark. So the universal mock theta function is the main influence on these expected values for large and small $|z|$, with appropriate choice of q .¹

¹For instance, take $q = e^{-1/kT}$ where k is Boltzmann's constant and T is the temperature of a system.

Conversely, if we write

$$\langle j_z \rangle_q =: \sum_{n=0}^{\infty} c_n q^n, \quad g_3(z^{-1}, q^{-1}) =: \sum_{n=0}^{\infty} \gamma_n q^n,$$

where the coefficients $c_n = c_n(z)$, $\gamma_n = \gamma_n(z)$ also depend on z , then the author proves an explicit combinatorial formula for the c_n in [39] using nested sums over subpartitions of n , viz.

$$(9) \quad c_n(z) = (1-z)^{-1} \sum_{\lambda \vdash n} \sum_{\delta | \lambda} \sum_{\epsilon | \delta} \sum_{\varphi | \epsilon} \mu(\lambda/\delta) z^{\text{crk}(\varphi)},$$

where, in the notations of [39], $\mu(\alpha) = 0$ if $\alpha \in \mathcal{P}$ has any part repeated and $= (-1)^{\ell(\alpha)}$ otherwise (a partition-theoretic version of the Möbius function first considered by Alladi [2, 39], where “ ℓ ” is the length of the partition), and if $\alpha | \beta$ then “ β/α ” denotes the partition obtained by deleting the parts of α from β .

With (9) in hand, it follows from Corollary 3 that the coefficients of $g_3(z^{-1}, q^{-1})$ satisfy the asymptotic

$$\gamma_n(z) \sim \begin{cases} z^{-1}(c_1 + c_2 + \dots + c_n) & \text{as } |z| \rightarrow \infty \\ z c_{n-1} & \text{as } |z| \rightarrow 0, \quad n \geq 1 \end{cases}$$

(which depends entirely on the growth of z , not n), as the coefficients enjoy the recursion

$$\gamma_n - \gamma_{n-1} \sim z^{-1} c_n \text{ for } |z| \gg 1.$$

Another observation: it is well known that if $\zeta_* \neq 1$ is a root of unity, then

$$(\zeta_* q; q)_{\infty} (\zeta_*^{-1} q; q)_{\infty}$$

is, up to multiplication by a rational power of q , a modular function [32]; but this product is the reciprocal of

$$(1 - \zeta_*) \cdot \langle j_z \rangle_q \Big|_{z=\zeta_*}.$$

This is another example of the intersection of the q -bracket with modularity phenomena, and at the same time gives a feeling for the obstruction to the inverted mock theta function’s sharing in this modularity at $z = \zeta_*$; for $g_3(z^{-1}, q^{-1})$ is not necessarily a dominating aspect of $\langle j_z \rangle_q$ for $z \neq 1$ near the unit circle, whereas the unimodal rank generating aspect $\tilde{U}(z, q)$ makes a more noticeable contribution, and the two pieces work together to produce modular behavior.

Going a little farther in this direction, there is a close relation between g_3 and the more general class of k -fold unimodal rank generating functions. Let us define the *rank generating function* $\tilde{U}_k(z, q)$ for unimodal sequences with a k -fold peak by the series

$$(10) \quad \tilde{U}_k(z, q) := \sum_{n=0}^{\infty} \frac{q^{kn}}{(zq; q)_n (z^{-1}q; q)_n},$$

where this identity follows directly from the combinatorial definition of $\tilde{U}_k$ [30].

Then it is not hard to find (see Theorem 1.1 of [38]) relations like

$$(11) \quad \frac{1}{(zq; q)_{\infty} (z^{-1}q; q)_{\infty}} = 2 - z - z^{-1} + (z + z^{-1})\tilde{U}_1(z, q) - \tilde{U}_2(z, q),$$

which of course is equal to $(1-z)\langle j_z \rangle_q$ and is modular for $z = \zeta_*$, up multiplication by a power of q . For example, noting that $z + z^{-1} = 0$ when $z = i$, then (11) yields

$$(12) \quad 2 - \tilde{U}_2(i, q) = (iq; q)_\infty^{-1}(-iq; q)_\infty^{-1} = (-q^2; q^2)_\infty^{-1},$$

where $(-q^2; q^2)_\infty$ is essentially a modular function.

At this point we can compare (11) to Theorem 2(i) to solve for $g_3(z^{-1}, q^{-1})$ in terms of $\tilde{U}_1, \tilde{U}_2$, but it is a little messy. However, it follows from a convenient rewriting of the right-hand side of Theorem 2(ii) using geometric series

$$\sum_{n=0}^{\infty} \frac{z}{(z; q)_{n+1}(z^{-1}q; q)_n} \left(\frac{z^{-1}q^{n+1}}{1 - z^{-1}q^{n+1}} \right) = \frac{z}{1-z} \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{z^{-k}q^{k(n+1)}}{(zq; q)_n(z^{-1}q; q)_n},$$

then swapping order of summation, that in fact $g_3(z^{-1}, q^{-1})$ can be written nicely in terms of the $\tilde{U}_k$.

Corollary 4. *For $|q| < 1 < |z|$, we have*

$$g_3(z^{-1}, q^{-1}) = \frac{z}{1-z} \sum_{k=1}^{\infty} \tilde{U}_k(z, q) z^{-k} q^k.$$

Thus the inverted universal mock theta function generates, in a sense, the sequence of rank generating functions for unimodal sequences with k -fold peaks, $k = 1, 2, 3, \dots$

Proof of Theorem 2. We begin by noting

$$\langle j_z \rangle_q = (z; q)_\infty^{-1}(z^{-1}q; q)_\infty^{-1} = \prod_{n=0}^{\infty} (1 - q^n(z + z^{-1}q - q^{n+1}))^{-1},$$

where in the final step we multiplied together the n th terms from each q -Pochhammer symbol. Thus we have

$$\prod_{n=0}^{\infty} (1 - q^n(z + z^{-1}q - q^{n+1}))^{-1} = 1 + \sum_{n=1}^{\infty} \frac{q^n(z + z^{-1}q - q^{n+1})}{\prod_{j=0}^{n-1} (1 - q^j(z + z^{-1}q - q^{j+1}))},$$

which can be shown by expanding the product on the left as the telescoping series

$$(13) \quad 1 + \sum_{n=1}^{\infty} \left(\frac{1}{\prod_{i=0}^n (1 - q^i(z + z^{-1}q - q^{i+1}))} - \frac{1}{\prod_{i=0}^{n-1} (1 - q^i(z + z^{-1}q - q^{i+1}))} \right)$$

with a little arithmetic (for more details see the proof of Theorem 1.1 (2) in [38]). Reorganizing the equations above, we arrive at the following relation.

Lemma 5. *For $|q| < 1, z \neq 0$, we have*

$$\langle j_z \rangle_q = 1 + (z + z^{-1}q) \sum_{n=1}^{\infty} \frac{q^n}{(z; q)_n(z^{-1}q; q)_n} - q \sum_{n=1}^{\infty} \frac{q^{2n}}{(z; q)_n(z^{-1}q; q)_n}.$$

We cannot help but notice how both series on the right-hand side of Lemma 5 resemble the right-hand summation of identity (6) for $\tilde{U}(z, q)$. This is not a coincidence; it follows right away from the simple observation

$$\tilde{U}(z, q) = \sum_{n=0}^{\infty} \frac{q^n}{(zq; q)_n(z^{-1}q; q)_n} = q^{-1}(1-z) \sum_{n=0}^{\infty} \frac{q^{n+1}(1 - z^{-1}q^{n+1})}{(z; q)_{n+1}(z^{-1}q; q)_{n+1}},$$

that $\tilde{U}$ splits off in a very similar fashion to $\langle j_z \rangle_q$ in Lemma 5, after taking into account $q \neq 0$:

$$(14) \quad \tilde{U}(z, q) = q^{-1}(1 - z) \sum_{n=1}^{\infty} \frac{q^n}{(z; q)_n (z^{-1}q; q)_n} - (zq)^{-1}(1 - z) \sum_{n=1}^{\infty} \frac{q^{2n}}{(z; q)_n (z^{-1}q; q)_n}$$

Comparing Lemma 5 and (14), plus a little bit of algebra, then gives

$$(15) \quad \langle j_z \rangle_q = 1 + [z(1 - q) + z^{-1}q] \sum_{n=1}^{\infty} \frac{q^n}{(z; q)_n (z^{-1}q; q)_n} + \frac{zq^2}{1 - z} \tilde{U}(z, q).$$

Now, to connect the remaining summation in (15) to the universal mock theta function g_3 , we apply a slightly clever factorization strategy in the q -Pochhammer symbols used by L. Rolin and the author in [37], eq. (4.1):

$$(16) \quad \begin{aligned} (z; q)_n (z^{-1}q; q)_n &= \prod_{j=0}^{n-1} [(-zq^j)(1 - z^{-1}(q^{-1})^j)] [(-z^{-1}q^{j+1})(1 - z(q^{-1})^{j+1})] \\ &= q^{n^2} (z^{-1}; q^{-1})_n (zq^{-1}; q^{-1})_n \end{aligned}$$

Thus

$$(17) \quad \sum_{n=1}^{\infty} \frac{q^n}{(z; q)_n (z^{-1}q; q)_n} = \sum_{n=1}^{\infty} \frac{q^n}{q^{n^2} (z^{-1}; q^{-1})_n (zq^{-1}; q^{-1})_n} = \sum_{n=1}^{\infty} \frac{(q^{-1})^{n(n-1)}}{(z^{-1}; q^{-1})_n (zq^{-1}; q^{-1})_n}.$$

The right-hand side of (17) is $g_3(z^{-1}, q^{-1})$, noting that it converges under the same conditions as the left side (being merely a term-wise rewriting), but with $q = 0$ omitted from the domain.

Remark. *Equivalently, identities like these result from the observation that*

$$(1 - zq^i)(1 - z^{-1}q^{-i})^{-1} = -zq^i.$$

Taking the product over $0 \leq i \leq n - 1$ gives

$$(z; q)_n (z^{-1}; q^{-1})_n^{-1} = (-1)^n z^n q^{n(n-1)/2}$$

and, proceeding in this manner, a variety of q -series summand forms can be produced (and inverted as above) by creative manipulation.

□

Remark. *We note in passing that, using (7.2) and (8.2) of Fine [18], Ch. 1, together with Theorem 2(ii), we can also write*

$$g_3(z^{-1}, q^{-1}) = (z^{-1}q; q)_{\infty}^{-1} (-z; q)_{\infty}^{-1} \sum_{n=0}^{\infty} (-1)^n z^{-2n} q^{\frac{n(n+1)}{2}} - \sum_{n=0}^{\infty} z^{-n+1} (z^{-1}; q)_n.$$

Recall that many modular forms arise as specializations of $j(z; q)$ (because $j(z; q)$ is essentially a Jacobi form [10]), and that $g_3(z, q)$ is the prototype for the class of mock modular forms that (to slightly abuse Ramanujan's words) "enter into mathematics as

beautifully” as the modular cases [23]. It is interesting that these important number-theoretic objects which are speculatively associated in the literature to opposite extremes of the universe—subatomic and supermassive—are themselves intertwined via the q -bracket from statistical physics, which applies to phenomena at every scale.

3. APPROACHING ROOTS OF UNITY RADially FROM WITHIN (AND WITHOUT)

One point that arises in Section 2 is that, apparently, the universal mock theta function g_3 can be written (at least formally) as a piecewise function

$$(18) \quad g_3(z, q) = \begin{cases} \sum_{n=1}^{\infty} \frac{q^{n(n-1)}}{(z; q)_n (z^{-1}q; q)_n} & \text{if } |q| < 1, \\ \sum_{n=1}^{\infty} \frac{(q^{-1})^n}{(z; q^{-1})_n (z^{-1}q^{-1}; q^{-1})_n} & \text{if } |q| > 1, \end{cases}$$

for q inside or outside the unit circle, respectively, and $z \neq 0$ or 1 . What of $g_3(z, q)$ for q lying on the circle? Generically one expects this question to be somewhat dicey.

To be precise in what follows, for ζ on the unit circle we define $g_3(z, \zeta)$ to mean the limit of $g_3(z, q)$ as $q \rightarrow \zeta$ radially from within (or without if the context allows), when the limit exists. Recalling the usual notation $\zeta_m := e^{2\pi i/m}$, it turns out that for $\zeta = \zeta_*$ an appropriate root of unity, $g_3(z, \zeta_*)$ is finite, both in value and length of the sum.

Theorem 6. *For $q = \zeta_m$ a primitive m th root of unity, $z \neq 0, 1$, or a rational power of ζ_m , and $z^m + z^{-m} \neq 1$, the odd-order universal mock theta function is given by the finite formula*

$$g_3(z, \zeta_m) = (1 - z^m - z^{-m})^{-1} \sum_{n=0}^{m-1} \zeta_m^n (z; \zeta_m)_n (z^{-1}\zeta_m; \zeta_m)_n.$$

Remark. Bringmann–Rolen [12] and Jang–Löbrich [26] have studied radial limits of universal mock theta functions from other perspectives.

Thus, under the right conditions, (18) together with Theorem 6 suggest $g_3(z, q)$ can, in a certain sense, “pass through” the unit circle at roots of unity (as a function of q following a radial path) into the complex plane beyond, and vice versa, while always remaining finite.

In the theory of quantum modular forms one encounters functions that exhibit similar behavior [10].

Definition 7. *Following Zagier [46], we say a function $f : \mathbb{P}^1(\mathbb{Q}) \setminus S \rightarrow \mathbb{C}$, for a discrete subset S , is a quantum modular form if $f(x) - f|_k \gamma(x) = h_\gamma(x)$ for a “suitably nice” function $h_\gamma(x)$, where $\gamma \in \Gamma$ a congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$, $|_k$ is the usual Petersson slash operator (see [32]), and “suitably nice” implies some pertinent analyticity condition, e.g. $\mathcal{C}_k, \mathcal{C}_\infty$, etc.*

These are functions that, in addition to being “almost” modular, generically “blow up” as q approaches the unit circle from within, but are finite when q radially approaches certain roots of unity or other isolated points, in which case the limiting values can be related to special values of L -functions [10]—and might even extend to the complex plane beyond the unit circle as in (18), a phenomenon called *renormalization* (see [35]). We see that g_3 exhibits this type of renormalization behavior.

Some mock theta functions are closely related to quantum modular forms. As we noted in Section 1, Ramanujan's mock theta function $f(q)$ (from eq. (1)) is, at even-order roots of unity, essentially a quantum modular form plus a modular form, through its relation to another rank generating function, the *rank generating function* $U(z, q)$ for *strongly unimodal sequences* [11, 19]:

$$(19) \quad U(z, q) := \sum_{n=0}^{\infty} q^{n+1} (-zq; q)_n (-z^{-1}q; q)_n$$

This $U(z, q)$ is a function that strikes deep: up to multiplication by rational powers of q , $U(i, q)$ is mock modular, $U(1, q)$ is mixed mock modular, and $U(-1, q)$ is a quantum modular form that can be completed to yield a weight $3/2$ non-holomorphic modular form [10].

Of course, U is the $k = 1$ case of the *rank generating function* $U_k(z, q)$ for *strongly unimodal sequences with k -fold peak*, given by

$$U_k(z, q) := \sum_{n=0}^{\infty} q^{k(n+1)} (-zq; q)_n (-z^{-1}q; q)_n.$$

Once again, we note the symmetry $U_k(z^{-1}, q) = U_k(z, q)$. As with $\tilde{U}_k$ in (11), we can find (see Theorem 2.8 of [38]) nice relations like

$$(20) \quad (zq; q)_{\infty} (z^{-1}q; q)_{\infty} = 1 - (z + z^{-1})U_1(z, q) + U_2(z, q),$$

which is modular for $z = \zeta_*$ a root of unity, up to multiplication by a power of q . For instance, at $z = i$, equation (20) gives

$$(21) \quad 1 + U_2(i, q) = (iq; q)_{\infty} (-iq; q)_{\infty} = (-q^2; q^2)_{\infty}.$$

Remark. Multiplying (12) and (21) leads to a nice pair of identities relating U_2 and $\tilde{U}_2$:

$$U_2(i, q) = \frac{1 - \tilde{U}_2(i, q)}{\tilde{U}_2(i, q) - 2}, \quad \tilde{U}_2(i, q) = \frac{1 + 2U_2(i, q)}{1 + U_2(i, q)}$$

Now, taking a similar approach to that in Section 2 with regard to $\tilde{U}_k$, we can find from Theorem 6, using an evaluation of $U_k(-z, q)$ at $q = \zeta_m$ much like the theorem²

$$(22) \quad U_k(-z, \zeta_m) = \frac{-1}{1 - z^m - z^{-m}} \sum_{n=0}^{m-1} \zeta_m^{k(n+1)} (z\zeta_m; \zeta_m)_n (z^{-1}\zeta_m; \zeta_m)_n,$$

that the universal mock theta function g_3 also connects to these rank generating functions U_k at roots of unity, through a similar relation to Corollary 4.

Corollary 8. For $|z| < 1$, we have

$$g_3(z, \zeta_m) = \frac{z - 1}{z} \sum_{k=1}^{\infty} U_k(-z, \zeta_m) z^k \zeta_m^{-k}.$$

How suggestive it is, in light of the relationship between $f(q)$ and $U(-1, q)$ [19], to see specializations of g_3 giving rise to both forms of k -fold unimodal rank generating functions in Corollaries 4 and 8.

²We note for $k = 1, z = 1, m$ even, that the summation in (22) appears in the right-hand side of (3).

Proof of Theorem 6 and Corollary 8. We start with an elementary observation. For an arbitrary q -series with coefficients d_n , then in the limit as q approaches an m th root of unity ζ_m radially from within the unit circle, we have

$$(23) \quad \lim_{q \rightarrow \zeta_m} \sum_{n=1}^{\infty} d_n q^n = \sum_{n=1}^m D_n \zeta_m^n \text{ where } D_n := \sum_{j=0}^{\infty} d_{n+mj},$$

so long as $\sum_j d_{n+mj}$ converges. The moral of this example: q -series *want* to be finite at roots of unity.

In a similar direction, Theorem 6 arises from the following very general lemma, which the author has spoken on at Emory University since 2012 and used for heuristics in projects with collaborators (such as [14, 37]), but has not published previously. It is really Lemma 9 below that is the pivotal result of Section 3; the applications to $g_3(z, \zeta_m)$ form an interesting exercise.

Lemma 9. *Suppose $\phi : \mathbb{Z}^+ \rightarrow \mathbb{C}$ is a periodic function of period $m \in \mathbb{Z}^+$, i.e., $\phi(r+mk) = \phi(r)$ for all $k \in \mathbb{Z}$. Define $f : \mathbb{Z}^+ \rightarrow \mathbb{C}$ by the product*

$$f(j) := \prod_{i=1}^j \phi(i),$$

and its summatory function $F(n)$ by $F(0) := 0$ and, for $n \geq 1$,

$$F(n) := \sum_{j=1}^n f(j), \quad F(\infty) := \lim_{n \rightarrow \infty} F(n) \text{ if the limit exists.}$$

Then the following statements are true:

(i) *For $0 \leq r < m$ we have*

$$F(mk + r) = \frac{1 - f(m)^k}{1 - f(m)} F(m) + f(m)^k F(r).$$

(ii) *For $|f(m)| < 1$ we have the finite formula*

$$F(\infty) = \frac{F(m)}{1 - f(m)}.$$

Proof of Lemma 9. First we observe that

$$(24) \quad f(mk) = \prod_{i=1}^{mk} \phi(i) = \left(\prod_{i=1}^m \phi(i) \right)^k = f(m)^k,$$

by the periodicity of ϕ . Then by the definition of $F(n)$ in Lemma 9 together with (24) we can rewrite

(25)

$$\begin{aligned}
F(mk + r) &= \sum_{j=1}^m f(j) + \sum_{j=m+1}^{2m} f(j) + \sum_{j=2m+1}^{3m} f(j) + \dots + \sum_{j=m(k-1)+1}^{mk} f(j) + \sum_{j=mk+1}^{mk+r} f(j) \\
&= \sum_{j=1}^m f(j) + f(m) \sum_{j=1}^m f(j) + f(m)^2 \sum_{j=1}^m f(j) + \dots + f(m)^{k-1} \sum_{j=1}^m f(j) + f(m)^k \sum_{j=1}^r f(j) \\
&= (1 + f(m) + f(m)^2 + \dots + f(m)^{k-1}) \sum_{j=1}^m f(j) + f(m)^k \sum_{j=1}^r f(j).
\end{aligned}$$

In the final expression above, recognizing the sum $1 + f(m) + f(m)^2 + \dots$ as a finite geometric series completes the proof of (i). If $|f(m)| < 1$, the infinite case gives (ii). $\square$

Remark. Then using Euler's continued fraction formula [17] we get another finite formula for $F(\infty)$, which holds for any convergent continued fraction of this shape with periodic coefficients:

$$F(\infty) = \frac{\phi(1)}{1 - \frac{\phi(2)}{1 + \phi(2) - \frac{\phi(3)}{1 + \phi(3) - \frac{\phi(4)}{1 + \phi(4) - \dots}}}} = \frac{1}{1 - f(m)} \left(\frac{\phi(1)}{1 - \frac{\phi(2)}{1 + \phi(2) - \frac{\phi(3)}{1 + \dots - \frac{\phi(m)}{1 + \phi(m)}}}} \right)$$

Many of the “quantum” considerations in this section also apply to continued fractions.

Clearly if we take ϕ to be sine, cosine, etc. in Lemma 9, we can produce a variety of trigonometric identities. More pertinently, if we replace $\phi(i)$ with $\tilde{\phi}(t, i) := t\phi(i)$, this $\tilde{\phi}$ also has period m ; then we see $\tilde{f}(j) := \prod_{i=1}^j \tilde{\phi}(t, i) = t^j f(j)$. Thus the summatory functions $\tilde{F}(n) = \tilde{F}(t, n)$ and $\tilde{F}(\infty) = \tilde{F}(t, \infty)$ represent a polynomial and a power series in t , respectively—which are, respectively, subject to (i) and (ii) of Lemma 9. Then for ϕ with period k and the product f as defined above, we get identities like

$$\sum_{n=1}^{\infty} f(n)t^n = \frac{1}{1 - f(k)t^k} \sum_{n=1}^k f(n)t^n.$$

(We could also take $\tilde{\phi}(t, i)$ equal to $t^i \phi(i)$ or $t^{2i} \phi(i)$ or $t^{2i-1} \phi(i)$, to lead to power series of other familiar shapes; however, such $\tilde{\phi}$ are not generally periodic.)

Thinking along these lines, if we set

$$\phi(i) = t \frac{(1 - a_1 q^{i-1})(1 - a_2 q^{i-1}) \dots (1 - a_r q^{i-1})}{(1 - b_1 q^{i-1})(1 - b_2 q^{i-1}) \dots (1 - b_s q^{i-1})}$$

for $a_*, b_* \in \mathbb{C}$, the product $f(j)$ becomes a quotient of q -Pochhammer symbols, producing the q -hypergeometric series

$$F(t, \infty) = {}_r F_s(a_1, \dots, a_r; b_1, \dots, b_s; t : q).$$

If $q \rightarrow \zeta_m$ an m th root of unity, then ϕ is also cyclic of period m , and in the radial limit ${}_r F_s(a_1, \dots, a_r; b_1, \dots, b_s; t : \zeta_m)$ is subject to Lemma 9(ii), so long as in the denominator $(1 - b_* \zeta_m^i) \neq 0$ for any i .

Remembering the “moral” of equation (23), then similar considerations apply to almost all q -series and Eulerian series, for $q = \zeta_m$ a root of unity that does not produce singularities. In particular, so long as the choice of z also does not lead to singularities, it is immediate from Lemma 9 by the definition (5) of g_3 that

$$(26) \quad \begin{aligned} g_3(z, \zeta_m) &= \frac{1}{1 - (z; \zeta_m)_m^{-1} (z^{-1} \zeta_m; \zeta_m)_m^{-1}} \sum_{n=1}^m \frac{\zeta_m^{n(n-1)}}{(z; \zeta_m)_n (z^{-1} \zeta_m; \zeta_m)_n} \\ &= \frac{2 - z^m - z^{-m}}{1 - z^m - z^{-m}} \sum_{n=1}^m \frac{\zeta_m^{n(n-1)}}{(z; \zeta_m)_n (z^{-1} \zeta_m; \zeta_m)_n}, \end{aligned}$$

where for the final equation we used the fact that

$$(X; \zeta_m)_m = 1 - X^m$$

in the leading factor. For a slightly simpler formula, we apply Lemma 9 to the identity for $g_3(z^{-1}; \zeta_m^{-1})$ in Theorem 2 instead, then take $z \mapsto z^{-1}$ and $\zeta_m \mapsto \zeta_m^{-1}$, to see

$$(27) \quad g_3(z, \zeta_m) = \frac{2 - z^m - z^{-m}}{1 - z^m - z^{-m}} \sum_{n=1}^m \frac{\zeta_m^{-n}}{(z^{-1}; \zeta_m^{-1})_n (z \zeta_m^{-1}; \zeta_m^{-1})_n}.$$

We note that the leading factor is symmetric under inversion of z .

Remark. Jang–Löbrich prove finite formulas similar to (26) and (27) for $g_3(z, \zeta_m)$ [26], by different methods.

A particularly lovely aspect of q -series such as these is that they transform into an infinite menagerie of shapes, limited only by the curiosity of the analyst. (For instance, see Fine [18] for a stunning exploration of q -hypergeometric series.) Then a form like g_3 might have a number of different finite formulas.

To derive Theorem 6, which is simpler than the preceding expressions for g_3 , we use another factorization strategy in the q -Pochhammer symbols. Again we exploit that

$$(X; \zeta_m)_m = 1 - X^m = (X; \zeta_m^{-1})_m;$$

thus for $0 \leq n \leq m$, since $\zeta_m^{-j} = \zeta_m^{m-j}$ we have

$$(28) \quad \begin{aligned} (X; \zeta_m^{-1})_n &= (1 - X)(1 - X \zeta_m^{m-1})(1 - X \zeta_m^{m-2}) \dots (1 - X \zeta_m^{m-(n-1)}) \\ &= \frac{(1 - X)(X; \zeta_m)_m}{(X; \zeta_m)_{m-n+1}} = \frac{(1 - X)(1 - X^m)}{(X; \zeta_m)_{m-n+1}}. \end{aligned}$$

Making the change of indices $n \mapsto m - n + 1$ in the summation in (27) then yields

$$\sum_{n=1}^m \frac{\zeta_m^{-(m-n+1)}}{(z^{-1}; \zeta_m^{-1})_{m-n+1} (z \zeta_m^{-1}; \zeta_m^{-1})_{m-n+1}} = \sum_{n=1}^m \frac{\zeta_m^{n-1} (z^{-1}; \zeta_m)_n (z \zeta_m^{-1}; \zeta_m)_n}{(1 - z^{-1})(1 - z \zeta_m^{-1})(2 - z^m - z^{-m})}.$$

Substituting this final expression back into (27), with a little algebra and adjusting of indices, gives Theorem 6.

To prove Corollary 8, we use geometric series, along with an order-of-summation swap and index change, to rewrite Theorem 6 in the form

$$\begin{aligned}
 (29) \quad g_3(z, \zeta_m) &= \frac{1-z}{(1-z^m-z^{-m})} \sum_{n=0}^{m-1} \zeta_m^n \frac{(z\zeta_m; \zeta_m)_n (z^{-1}\zeta_m; \zeta_m)_n}{1-z\zeta_m^n} \\
 &= \frac{z^{-1}(1-z)}{z(1-z^m-z^{-m})} \sum_{k=1}^{\infty} z^k \zeta_m^{-k} \sum_{n=0}^{m-1} \zeta_m^{k(n+1)} (\zeta_m; \zeta_m)_n (z^{-1}\zeta_m; \zeta_m)_n.
 \end{aligned}$$

Comparing this with the formula (22) for $U(-z, \zeta_m)$, which follows easily from Lemma 9, gives the corollary. (The sum on the right might be simplified further using (23).)

Remark. *Convergence in these formulas depends on one's choice of substitutions; for a particular choice, careful analysis may be required to show boundedness as q approaches the natural boundary of a q -series (see Watson [44] for examples).*

□

We note that a slight variation on the proof above leads to finite formulas at applicable roots of unity for the even-order universal mock theta function $g_2(z, q)$ of Gordon–McIntosh [21] as well, by an alternative approach to that of Bringmann–Rolen [12]. Using transformations from Andrews [3], Fine [18], and other authors, still simpler formulas might be found for particular specializations of g_3 at roots of unity. We demonstrate this point below.

Example 10. *The radial limit of $f(q)$ at ζ_m an odd-order root of unity is equal to*

$$f(\zeta_m) = 1 - \frac{2}{3} \sum_{n=1}^m (-1)^n \zeta_m^{-(n+1)} (-\zeta_m^{-1}; \zeta_m^{-1})_n.$$

Proof of Example 10. By (26.22) in [18], Ch. 3, Ramanujan's mock theta function $f(q)$ defined in (1) can be rewritten

$$(30) \quad f(q) = 1 - \sum_{n=1}^{\infty} \frac{(-1)^n q^n}{(-q; q)_n}.$$

To show the summation on the right is bounded as q approaches an odd-order root of unity radially, we exactly follow the steps of Watson's analysis of the mock theta function $f_0(q)$ in [44], Sec. 6. In Watson's nomenclature, take $M = 2$, N odd, to write $q = e^{2\pi i/N} = \zeta_N$. Then by replacing $q^{(nN+m)^2}$ with $(-1)^{nN+m} q^{nN+m}$ in the numerators of the $n \geq 1$ terms of the series $f_0(q)$ (we note that Watson's m is not the same as the subscript of ζ_m we use throughout this paper, which corresponds to N in this proof), one sees

$$\left| 1 - \sum_{n=1}^{\infty} \frac{(-1)^n q^n}{(-q; q)_n} \right| \leq 2 \sum_{n=0}^{N-1} \left| \frac{q^n}{(-q; q)_n} \right| < \infty$$

when $q = \rho \zeta_N$ with $0 \leq \rho \leq 1$. To see the value the series converges to, consider the $(Nk + r)$ th partial sum, with $r < N$, of the right-hand side of (30) as $\rho \rightarrow 1^-$, in light of Lemma 9 (i). In fact, as $|(-1)^N \zeta_N^N / (-\zeta_N, \zeta_N)_N| = 1/2 < 1$, then part (ii) of Lemma 9

applies as $Nk + r \rightarrow \infty$ and (also taking into account that $f(\rho\zeta_N)$ converges uniformly for $\rho < 1$) we may write

$$(31) \quad \lim_{\rho \rightarrow 1^-} \left(1 - \sum_{n=1}^{\infty} \frac{(-1)^n (\rho\zeta_N)^n}{(-\rho\zeta_N; \rho\zeta_N)_n} \right) = 1 - \frac{2}{3} \sum_{n=1}^N \frac{(-1)^n \zeta_N^n}{(-\zeta_N; \zeta_N)_n}.$$

Now, observe that (28) gives

$$(32) \quad (-\zeta_N; \zeta_N)_n^{-1} = (-\zeta_N^{-1}; \zeta_N^{-1})_{N-n-1}.$$

Applying (32) to the right-hand side of (31), then making the change $n \mapsto N - n - 1$ in the indices, we arrive at the desired result. $\square$

Continuing in this fashion, we can find a formula for this radial limit that is even easier to compute.

Example 11. For ζ_m an odd-order root of unity we have

$$f(\zeta_m) = \frac{4}{3} \sum_{n=1}^m (-1)^n (-\zeta_m^{-1}; \zeta_m^{-1})_n.$$

Proof of Example 11. Here we use only finite sums, so we do not need to justify convergence. Let us define an auxiliary series

$$g(\zeta_m) = \frac{2}{3} \sum_{n=1}^m (-1)^n (-\zeta_m^{-1}; \zeta_m^{-1})_n;$$

then using Example 10, with a little arithmetic and adjusting of indices, gives

$$f(\zeta_m) - g(\zeta_m) = 1 - \frac{2}{3} \sum_{n=1}^m (-1)^n (-\zeta_m^{-1}; \zeta_m^{-1})_{n+1} = g(\zeta_m).$$

Comparing the left- and right-hand sides above implies our claim. $\square$

Indeed, by the considerations here we can find both finite formulas at roots of unity, and inverted versions using factorizations such as in (16) leading to piecewise forms such as (18), for a great many q -hypergeometric series. Whether or not they enjoy modularity properties, these can display very interesting behaviors, emerging outside the unit circle radially from an entirely different point ζ_m^{-1} than the point on the circle ζ_m approached from within, and likewise when entering the circle radially at roots of unity from without. Moreover, the map inside the unit circle in the variable q looks like an “upside-down hyperbolic mirror-image” of the function’s behavior on the outside. (Taking $q \mapsto \bar{q}$ in either the $|q| < 1$ or $|q| > 1$ piece of (18) turns the map “right-side up”, but at the expense of holomorphicity.³)

This imagery reminds us of depictions of white holes and wormholes in science fiction. Do there exist “points-of-exit” (and entry) analogous in some way to roots of unity, at the event horizon of a black hole? Is there a mirror-image universe contained within? We don’t take these fantastical analogies too seriously, yet one is led to wonder: how deep is the connection between q -series and phenomena at nature’s fringe?

³As Tyler Smith, Emory University Department of Physics, noted to the author [40].

ACKNOWLEDGEMENTS

I am thankful to George Andrews for extremely helpful comments pertaining to Examples 10 and 11, and for suggesting the Watson references; to Jeremy Lovejoy for useful correspondence regarding unimodal rank generating functions; to Larry Rolin for offering revisions that strengthened this work; and to Krishnaswami Alladi, Olivia Beckwith, David Borthwick, Amanda Clemm, John Duncan, Amanda Folsom, Marie Jameson, Benjamin Phelan, Robert Rhoades, Tyler Smith, and my Ph.D. advisor, Ken Ono, for discussions that informed my study.

REFERENCES

- [1] B. P. Abbott, et al., *Observation of gravitational waves from a binary black hole merger*, Physical review letters **116.6** (2016): 061102.
- [2] K. Alladi, Private communication, December 22, 2015.
- [3] G. E. Andrews, *The theory of partitions*, no. 2, Cambridge University Press, 1998.
- [4] G. E. Andrews, *An introduction to Ramanujan's "lost" notebook*, The American Mathematical Monthly **86.2** (1979): 89-108.
- [5] G. E. Andrews and F. G. Garvan, *Dyson's crank of a partition*, Bull. Am. Math. Soc. **18.2** (1988).
- [6] T. M. Apostol, *Modular functions and Dirichlet series in number theory*, Vol. 41, Springer Science and Business Media, 2012.
- [7] B. C. Berndt, *Number theory in the spirit of Ramanujan*, Vol. 34. American Mathematical Soc., 2006.
- [8] S. Bloch and A. Okounkov, *The character of the infinite wedge representation*, Adv. Math. **149** (2000), no. 1, 1-60.
- [9] K. Bringmann, A. Folsom and R. C. Rhoades, *Unimodal sequences and "strange" functions: a family of quantum modular forms*, Pacific Journal of Mathematics **274.1** (2015): 1-25.
- [10] K. Bringmann, A. Folsom, K. Ono and L. Rolin, *Harmonic Maass forms and mock modular forms: Theory and applications*, American Mathematical Society, Providence, RI, In preparation.
- [11] J. Bryson, K. Ono, S. Pitman and R. C. Rhoades, *Unimodal sequences and quantum and mock modular forms*, Proceedings of the National Academy of Sciences **109.40** (2012): 16063-16067.
- [12] K. Bringmann and L. Rolin, *Radial limits of mock theta functions*, Research in the Mathematical Sciences, **2: 17** (2015).
- [13] P. J. Cameron, *Combinatorics: Topics, Techniques, Algorithms*, Cambridge University (1994).
- [14] A. Clemm and R. Schneider, *Results from a computational study of cyclotomic phenomena in the mock theta function $f(q)$* , Preprint (2016).
- [15] A. Dabholkar, S. Murthy and D. Zagier, *Quantum black holes, wall crossing, and mock modular forms*, arXiv preprint arXiv:1208.4074 (2012).
- [16] J. F. R. Duncan, M. J. Griffin, and K. Ono, *Moonshine*, Research in the Mathematical Sciences **2.1** (2015): 1-57.
- [17] L. Euler, *An essay on continued fractions*, Mathematical Systems Theory **18.4** (1985): 295-328.
- [18] N. J. Fine, *Basic hypergeometric series and applications*, With a foreword by George E. Andrews, Mathematical Surveys and Monographs, 27. American Mathematical Society, Providence, RI, 1988.
- [19] A. Folsom, K. Ono and R. C. Rhoades, *Mock theta functions and quantum modular forms*, Forum of Mathematics, Pi. Vol. 1. Cambridge University Press, 2013.
- [20] G. Gamow, *Thirty years that shook physics: The story of quantum theory*, Courier Corporation, 1966.
- [21] B. Gordon and R. J. McIntosh, *A survey of classical mock theta functions*, Partitions, q -series, and modular forms, 95144, Dev. Math., 23, Springer, New York, 2012.
- [22] M. Griffin, M. Jameson and S. Trebat-Leder, *On p -adic modular forms and the Bloch-Okounkov theorem*, arXiv preprint arXiv:1509.07161 (2015).

- [23] G. H. Hardy, *Ramanujan: Twelve lectures on subjects suggested by his life and work*, Chelsea Pub. Co., New York (1959).
- [24] D. R. Hickerson and E. T. Mortenson, *Hecke-type double sums, Appell-Lerch sums, and mock theta functions, I*, Proceedings of the London Mathematical Society (2014): pdu007.
- [25] W. Isaacson, *How Einstein reinvented reality*, Scientific American **313**(2015): 38-45.
- [26] M. Jang and S. L  brich, *Radial limits of the universal mock theta function g_3* , arXiv preprint arXiv:1504.05365 (2015).
- [27] B. Kim and J. Lovejoy, *The rank of a unimodal sequence and a partial theta identity of Ramanujan*, International Journal of Number Theory **10.04** (2014): 1081-1098.
- [28] B. Kim and J. Lovejoy, *Ramanujan-type partial theta identities and rank differences for special unimodal sequences*, Annals of Combinatorics **19.4** (2015): 705-733.
- [29] K. Knopp, *Theory of functions*, Vol. 1, Courier Corporation, 1996.
- [30] J. Lovejoy, Private communication, August 3, 2016.
- [31] K. Ono, *Unearthing the visions of a master: harmonic Maass forms and number theory*, Current developments in mathematics 2008 (2009): 347-454.
- [32] K. Ono, *The web of modularity: arithmetic of the coefficients of modular forms and q -series*, volume 102 of CBMS Regional Conference Series in Mathematics, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2004.
- [33] B. Phelan and R. Schneider, *Encounter with the infinite*, The Believer **13:1** (2015).
- [34] S. Ramanujan, *Collected papers*, Cambridge University Press, Cambridge, 1927; reprinted by Chelsea, New York, 1962; reprinted by the American Mathematical Society, Providence, 2000.
- [35] Y. Li, H. T. Ngo and R. C. Rhoades, *Renormalization and quantum modular forms, part I: Maass wave forms*, arXiv preprint arXiv:1311.3043 (2013).
- [36] R. Schneider, Uncovering Ramanujans "Lost" Notebook: an oral history, The Ramanujan Journal **29.1-3** (2012): 3-24.
- [37] L. Rolin and R. Schneider, *A "strange" vector-valued quantum modular form*, Archiv der Mathematik **101.1** (2013): 43-52.
- [38] R. Schneider, *Partition zeta functions*, Research in Number Theory **2: 9** (2016).
- [39] R. Schneider, *Arithmetic of partitions and the q -bracket operator*, Proc. of the Am. Math. Soc. (To appear).
- [40] T. Smith, Private communication (September 9, 2016).
- [41] J. Steinhauer, *Observation of quantum Hawking radiation and its entanglement in an analogue black hole*, Nature Physics (2016).
- [42] S. O. Warnaar, *Partial theta functions. I. Beyond the lost notebook*, Proceedings of the London Mathematical Society **87.2** (2003): 363-395.
- [43] G. N. Watson, *The final problem: an account of the mock theta functions*, Journal of the London Mathematical Society **1.1** (1936): 55-80.
- [44] G. N. Watson, *Mock theta functions (2)*, Proc. London Math. Soc. (2), **42** (1937), 274-304.
- [45] D. Zagier, *Partitions, quasimodular forms, and the Bloch-Okounkov theorem*, The Ramanujan Journal (2015): 1-24.
- [46] D. Zagier, *Quantum modular forms*, Quanta of maths **11** (2010): 659-675.
- [47] S. Zwegers, *Mock theta functions*, arXiv preprint arXiv:0807.4834 (2008).

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
 EMORY UNIVERSITY
 400 DOWMAN DR., W401
 ATLANTA, GEORGIA 30322
E-mail address: robert.schneider@emory.edu